

Anti-Protein Kinase C Eta Recombinant Rabbit Monoclonal Antibody



Catalog #: 2576

Aliases

PRKCH; Protein Kinase C Eta; PKC-L; PKCL; PRKCL; Protein Kinase C Eta Type; Protein UPEP; EC 2.7.11.13; NPKC-Eta; UORF2; PRKCH Upstream Open Reading Frame 2; Protein Kinase C, Eta; EC 2.7.11

Background

Gene Name: PRKCH

NCBI Gene Entry: [5583](#)

UniProt Entry: [P24723](#)

Application Information

Molecular Weight: Predicted, 78 kDa; observed, 78 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 24GB1480

Species Reactivity: Human, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

Immunogen

A synthesized peptide derived from human PKC eta

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions

Western Blotting (WB): 1:1,000-1:5,000

Flow Cytometry (FCM): 1:2,000

Immunocytochemistry (IC): 1:100-1:1,000

Note: This product is for research use only.

SUPPORT


SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041


WWW.GENUINBIOTECH.COM

Validation Data



Copyright ©2024 Genuin Biotechnologies LLC

Flow cytometric analysis of Protein kinase C eta expression in HepG2 cells using anti-Protein kinase C eta antibody (Cat#2576, 1:2,000). Green, isotype control; red, Protein kinase C eta.



Copyright ©2024 Genuin Biotechnologies LLC

Immunocytochemical staining of HepG2 cells with anti-Protein kinase C eta antibody (Cat#2576, 1:1,000). Nuclei were stained blue with DAPI; Protein kinase C eta was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 μm.


SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM



Copyright © 2024 Genuin Biotechnologies LLC

Western blotting analysis using anti-Protein kinase C eta antibody (Cat#2576). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Protein kinase C eta antibody (Cat#2576, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQ™ ECL Substrate Kit (Cat#226).