

Anti-HTR Recombinant Rabbit Monoclonal Antibody



Catalog #: 2979

Aliases

HTR; 5-Hydroxytryptamine Receptor 2C; 5-HT2C; 5HTR2C; HTR1C; 5-Hydroxytryptamine (Serotonin) Receptor 2C, G Protein-Coupled; 5-Hydroxytryptamine Receptor 1C; 5-HTR2C; 5-HT1C; 5-Hydroxytryptamine (Serotonin) Receptor 2C; Serotonin 5-HT-1C Receptor; Serotonin 5-HT-2C Receptor; Serotonin Receptor 2C; 5-HT-2C; 5-HT-1C

Background

Gene Name: HTR2C

NCBI Gene Entry: [3358](#)

UniProt Entry: [P28335](#)

Application Information

Molecular Weight: Predicted, 52 kDa; observed, 73 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 24GB3460

Species Reactivity: Human, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

Immunogen

A synthesized peptide derived from human 5HT2C Receptor

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions

Western Blotting (WB): 1:1,000-1:5,000

Flow Cytometry (FCM): 1:2,000

Immunocytochemistry (IC): 1:100-1:1,000

SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041


WWW.GENUINBIOTECH.COM

Anti-HTR Recombinant Rabbit Monoclonal Antibody

PAGE 2

Note: This product is for research use only.


Validation Data



Copyright ©2024 Genuin Biotechnologies LLC

5-hydroxytryptamine receptor 2C-Alexa Fluor® 647

Flow cytometric analysis of 5-hydroxytryptamine receptor 2C expression in HepG2 cells using anti-5-hydroxytryptamine receptor 2C antibody (Cat#2979, 1:2,000). Green, isotype control; red, 5-hydroxytryptamine receptor 2C.



Copyright ©2024 Genuin Biotechnologies LLC

Immunocytochemical staining of HepG2 cells with anti-5-hydroxytryptamine receptor 2C

SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS


SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM

Anti-HTR Recombinant Rabbit Monoclonal Antibody

PAGE 3

antibody (Cat#2979, 1:1,000) . Nuclei were stained blue with DAPI; 5-hydroxytryptamine receptor 2C was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar, 20 μ m.



Western blotting analysis using anti-5-hydroxytryptamine receptor 2C antibody (Cat#2979). Total cell lysates (30 μ g) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-5-hydroxytryptamine receptor 2C antibody (Cat#2979, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQ™ ECL Substrate Kit (Cat#226).

SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM