# **Anti-NME2 Recombinant Rabbit Monoclonal Antibody**



**Catalog #: 3033** 

#### **Aliases**

NME2; NME/NM23 Nucleoside Diphosphate Kinase 2; NM23-H2; NDPKB; Non-Metastatic Cells 2, Protein (NM23B) Expressed In; C-Myc Purine-Binding Transcription Factor PUF; Nucleoside Diphosphate Kinase B; Histidine Protein Kinase NDKB; NDP Kinase B; EC 2.7.4.6; NM23B; Epididymis Secretory Sperm Binding Protein Li 155an; Non-Metastatic Cells 2, Protein (NM23) Expressed In; Nucleotide Diphosphate Kinase B; C-Myc Transcription Factor; HEL-S-155an; EC 2.7.13.3; Nm23-H2; NDPK-B; NDK B; NDKB; PUF

# **Background**

Gene Name: NME2 NCBI Gene Entry: 4831 UniProt Entry: P22392

# **Application Information**

Molecular Weight: Predicted, 17 kDa; observed, 17 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 24GB3730

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

#### Immunogen

A synthesized peptide derived from human NME2

### **Isotype**

Rabbit IgG

### **Storage Buffer**

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

# **Storage**

Store at -20 °C for one year.

## **Recommended Dilutions**


Western Blotting (WB): 1:1,000-1:5,000

Flow Cytometry (FCM): 1:2,000

Immunocytochemistry (IC): 1:100-1:1,000


**Note:** This product is for research use only.

#### **Validation Data**




Copyright @2024 Genuin Biotechnologies LLC

Flow cytometric analysis of NME2 expression in HepG2 cells using anti-NME2 antibody (Cat#3033, 1:2,000). Green, isotype control; red, NME2.



Immunocytochemical staining of HepG2 cells with anti-NME2 antibody (Cat#3033, 1:1,000) . Nuclei were stained blue with DAPI; NME2 was stained magenta with Alexa Fluor® 647. Images were taken using leica stellaris 5. Protein abundance based on laser intensity and smart gain: Medium. Scale bar:  $20~\mu m$ .

# **Anti-NME2 Recombinant Rabbit Monoclonal Antibody**



Western blotting analysis using anti-NME2 antibody (Cat#3033). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-NME2 antibody (Cat#3033, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQ<sup>TM</sup> ECL Substrate Kit (Cat#226).