Anti-Phospho-CBL (S669) Recombinant Rabbit Monoclonal Antibody



Catalog #: 3061

Aliases

CBL; Cbl Proto-Oncogene; RNF55; C-Cbl; CBL2; Cas-Br-M (Murine) Ecotropic Retroviral Transforming Sequence; Cbl Proto-Oncogene, E3 Ubiquitin Protein Ligase; Casitas B-Lineage Lymphoma Proto-Oncogene; RING-Type E3 Ubiquitin Transferase CBL; E3 Ubiquitin-Protein Ligase CBL; Signal Transduction Protein CBL; RING Finger Protein 55; Proto-Oncogene C-Cbl; Oncogene CBL2; Fragile Site, Folic Acid Type, Rare, Fra(11)(Q23.3); EC 2.3.2.27; EC 6.3.2; FRA11B; NSLL

Background

Gene Name: CBL NCBI Gene Entry: 867 UniProt Entry: P22681

Application Information

Molecular Weight: Predicted, 100 kDa; observed, 120 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 24GB3870

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

Immunogen

A synthesized peptide derived from human Phospho-CBL (S669)

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions

Western Blotting (WB): 1:1,000-1:5,000


Flow Cytometry (FCM): 1:2,000

Immunocytochemistry (IC): 1:100-1:1,000


TEL: +1-540-855-7041

Note: This product is for research use only.

Validation Data




Flow cytometric analysis of phospho-CBL (S669) expression in HepG2 cells using anti-phospho-CBL (S669) antibody (Cat#3061, 1:2,000). Green, isotype control; red, phospho-CBL (S669).



Immunocytochemical staining of HepG2 cells with anti-Phospho-CBL (S669) antibody (Cat#3061, 1:1,000). Nuclei were stained blue with DAPI; Phospho-CBL (S669) was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 µm.

Anti-Phospho-CBL (S669) Recombinant Rabbit Monoclonal Antibody



Western blotting analysis using anti-phospho-CBL (S669) antibody (Cat#3061). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-phospho-CBL (S669) antibody (Cat#3061, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQTM ECL Substrate Kit (Cat#226).