# **Anti-Sequestosome 1 Recombinant Rabbit Monoclonal Antibody**



**Catalog #: 3133** 

#### **Aliases**

SQSTM1; Sequestosome 1; P60; P62B; A170; OSIL; P62; Phosphotyrosine-Independent Ligand For The Lck SH2 Domain Of 62 KDa; EBI3-Associated Protein Of 60 KDa; Oxidative Stress Induced Like; Ubiquitin-Binding Protein P62; Autophagy Receptor P62; Sequestosome-1; EBIAP; PDB3; Phosphotyrosine Independent Ligand For The Lck SH2 Domain P62; EBI3-Associated Protein P60; Paget Disease Of Bone 3; FTDALS; NADGP; DMRV; ZIP3; ORCA

## **Background**

Gene Name: SQSTM1 NCBI Gene Entry: 8878 UniProt Entry: Q13501

## **Application Information**

Molecular Weight: Predicted, 48 kDa; observed, 62 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 24GB4225

Species Reactivity: Human

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

#### Immunogen

A synthesized peptide derived from human SQSTM1 / p62

## **Isotype**

Rabbit IgG

### **Storage Buffer**

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

### **Storage**

Store at -20 °C for one year.

#### **Recommended Dilutions**

Western Blotting (WB): 1:1,000-1:5,000


Flow Cytometry (FCM): 1:2,000

Immunocytochemistry (IC): 1:100-1:1,000


Copyright @2024 Genuin Biotechnologies LLC

Note: This product is for research use only.

#### **Validation Data**




Flow cytometric analysis of p62/SQSTM1 expression in HepG2 cells using anti-p62/SQSTM1 antibody (Cat#3133, 1:2,000). Green, isotype control; red, p62/SQSTM1.



# **Anti-Sequestosome 1 Recombinant Rabbit Monoclonal Antibody**

Immunocytochemical staining of HepG2 cells with anti-p62/SQSTM1 antibody (Cat#3133, 1:1,000). Nuclei were stained blue with DAPI; p62/SQSTM1 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar: 20 μm.



Western blotting analysis using anti-p62/SQSTM1 antibody (Cat#3133). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-p62/SQSTM1 antibody (Cat#3133, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQ<sup>TM</sup> ECL Substrate Kit (Cat#226).