Anti-Peptidylprolyl isomerase F Recombinant Rabbit Monoclonal Antibody



Catalog #: 4365

Aliases

PPIF; Peptidylprolyl Isomerase F; Cyclophilin D; CypD; Peptidyl-Prolyl Cis-Trans Isomerase F, Mitochondrial; Mitochondrial Cyclophilin; Cyclophilin F; Rotamase F; EC 5.2.1.8; PPIase F; Cyp-D; CyP-M; CYP3; Peptidyl-Prolyl Cis-Trans Isomerase, Mitochondrial; Peptidylprolyl Isomerase F (Cyclophilin F; Cyclophilin 3; HCyP3; CYP-D; HCYP3; CyP-D; CYPD

Background

Gene Name: PPIF

NCBI Gene Entry: 10105 UniProt Entry: P30405

Application Information

Molecular Weight: Predicted, 22 kDa; observed, 18 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 24GB10855

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

Immunogen

A synthesized peptide derived from human Cyclophilin F

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions

Western Blotting (WB): 1:1,000-1:5,000


Flow Cytometry (FCM): 1:2,000

Immunocytochemistry (IC): 1:100-1:1,000

Anti-Peptidylprolyl isomerase F Recombinant Rabbit Monoclonal Antibody

Note: This product is for research use only.

Validation Data




Copyright ©2024 Genuin Biotechnologies LLC


Peptidylprolyl isomerase F-Alexa Fluor® 647

Flow cytometric analysis of peptidylprolyl isomerase F expression in C2C12 cells using antipeptidylprolyl isomerase F antibody (Cat#4365, 1:2,000). Green, isotype control; red, peptidylprolyl isomerase F.

Anti-Peptidylprolyl isomerase F Recombinant Rabbit Monoclonal Antibody



Immunocytochemical staining of C2C12 cells with anti-Peptidylprolyl isomerase F antibody (Cat#4365, 1:1,000). Nuclei were stained blue with DAPI; Peptidylprolyl isomerase F was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar, 20 μm.



Western blotting analysis using anti-peptidylprolyl isomerase F antibody (Cat#4365). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-peptidylprolyl isomerase F antibody (Cat#4365, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using NaQTM ECL Substrate Kit (Cat#716).