## **Anti-PPM1G Recombinant Rabbit Monoclonal Antibody**



### **Catalog #: 4422**

#### **Aliases**

PPM1G; Protein Phosphatase, Mg2+/Mn2+ Dependent 1G; PP2CG; Protein Phosphatase Magnesium-Dependent 1 Gamma; Protein Phosphatase 1G; Protein Phosphatase 1C; EC 3.1.3.16; PP2C-Gamma; PP2CGAMMA; Protein Phosphatase 1G (Formerly 2C), Magnesium-Dependent, Gamma Isoform; Protein Phosphatase, Mg2+/Mn2+ Dependent, 1G; Protein Phosphatase 2C, Gamma Isoform; Protein Phosphatase 2C Isoform Gamma; PP2C, Gamma; PP2Cgamma; PPP2CG; PPM1C

### **Background**

Gene Name: PPM1G NCBI Gene Entry: 5496 UniProt Entry: O15355

## **Application Information**

Molecular Weight: Predicted, 59 kDa; observed, 72 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 24GB11140

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

#### **Immunogen**

A synthesized peptide derived from human PPM1G

### **Isotype**

Rabbit IgG

#### **Storage Buffer**

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

#### **Storage**

Store at -20 °C for one year.

#### **Recommended Dilutions**


Western Blotting (WB): 1:1,000-1:5,000

Flow Cytometry (FCM): 1:2,000

Immunocytochemistry (IC): 1:100-1:1,000

**Note:** This product is for research use only.


#### **Validation Data**



Flow cytometric analysis of PPM1G expression in HepG2 cells using anti-PPM1G antibody (Cat#4422, 1:2,000). Green, isotype control; red, PPM1G.


Copyright ©2024 Genuin Biotechnologies LLC

# **Anti-PPM1G Recombinant Rabbit Monoclonal Antibody**



Immunocytochemical staining of HepG2 cells with anti-PPM1G antibody (Cat#4422, 1:1,000). Nuclei were stained blue with DAPI; PPM1G was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar,  $20~\mu m$ .

# **Anti-PPM1G Recombinant Rabbit Monoclonal Antibody**



Western blotting analysis using anti-PPM1G antibody (Cat#4422). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-PPM1G antibody (Cat#4422, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using NaQ<sup>TM</sup> ECL Substrate Kit (Cat#716).