#### **Anti-Phospho-PKC alpha (Thr638) Rabbit Monoclonal Antibody**



**Catalog #: 4684** 

#### **Aliases**

PRKCA; Protein Kinase C Alpha; PKCA; Protein Kinase C Alpha Type; EC 2.7.11.13; PKC-Alpha; PRKACA; PKC-A; Protein Kinase C, Alpha; Aging-Associated Gene 6; EC 2.7.11; PKCalpha; PKCI+/-; PKCα; AAG6

## **Background**

Gene Name: PRKCA NCBI Gene Entry: 5578 UniProt Entry: P17252

## **Application Information**

Molecular Weight: Predicted, 77 kDa; observed, 74 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 24GB12430

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

#### **Immunogen**

A synthesized peptide derived from human Phospho-PKC alpha (Thr638)

#### **Isotype**

Rabbit IgG

## **Storage Buffer**

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

### **Storage**

Store at -20 °C for one year.

#### **Recommended Dilutions**


Western Blotting (WB): 1:1,000-1:5,000

Flow Cytometry (FCM): 1:2,000

Immunocytochemistry (IC): 1:100-1:1,000

**Note:** This product is for research use only.


#### **Validation Data**



Copyright @2024 Genuin Biotechnologies LLC

## Phospho-PKC alpha (Thr638)-Alexa Fluor® 647


Flow cytometric analysis of phospho-PKC alpha (Thr638) expression in HepG2 cells using anti-phospho-PKC alpha (Thr638) antibody (Cat#4684, 1:2,000). Green, isotype control; red, phospho-PKC alpha (Thr638).



Immunocytochemical staining of HepG2 cells with anti-Phospho-PKC alpha (Thr638) antibody

# Anti-Phospho-PKC alpha (Thr638) Rabbit Monoclonal Antibody

(Cat#4684, 1:1,000). Nuclei were stained blue with DAPI; Phospho-PKC alpha (Thr638) was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar, 20 µm.



Western blotting analysis using anti-phospho-PKC alpha (Thr638) antibody (Cat#4684). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-phospho-PKC alpha (Thr638) antibody (Cat#4684, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using NaQ<sup>TM</sup> ECL Substrate Kit (Cat#716).