Anti-GCLC Rabbit Monoclonal Antibody



Catalog #: 4718

Aliases

GCLC; Glutamate-Cysteine Ligase Catalytic Subunit; GLCLC; GLCL; GCS; Glutamate--Cysteine Ligase Catalytic Subunit; Gamma-Glutamylcysteine Synthetase; GCS Heavy Chain; EC 6.3.2.2; Gamma-ECS; Glutamate-Cysteine Ligase, Catalytic Subunit; GCL

Background

Gene Name: GCLC NCBI Gene Entry: 2729 UniProt Entry: P48506

Application Information

Molecular Weight: Predicted, 61 kDa; observed, 73 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 24GB12600

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

Immunogen

Recombinant protein of human GCLC

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions


Western Blotting (WB): 1:1,000-1:5,000

Flow Cytometry (FCM): 1:2,000


Immunocytochemistry (IC): 1:100-1:1,000

Note: This product is for research use only.

Validation Data




Flow cytometric analysis of GCLC expression in HepG2 cells using anti-GCLC antibody (Cat#4718, 1:2,000). Green, isotype control; red, GCLC.



Immunocytochemical staining of HepG2 cells with anti-GCLC antibody (Cat#4718, 1:1,000). Nuclei were stained blue with DAPI; GCLC was stained magenta with Alexa Fluor® 647. Images

were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar, $20~\mu m$.



Western blotting analysis using anti-GCLC antibody (Cat#4718). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-GCLC antibody (Cat#4718, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQTM ECL Substrate Kit (Cat#226).