Anti-Kv1.3 Rabbit Polyclonal Antibody



Catalog #: 50440

Aliases

HGK5; Potassium voltage-gated channel subfamily A member 3; HGK5; HLK3; HPCN3; Voltage-gated K(+) channel HuKIII; Voltage-gated potassium channel subunit Kv1.3

Background

Gene Name: KCNA3 NCBI Gene Entry: 3738 UniProt Entry: P22001

Application Information

Molecular Weight: Predicted, 63 kDa; observed, 63 kDa

Clonality: Rabbit polyclonal antibody

Species Reactivity: Human, mouse, rat, bovine, chicken, pig, rabbit

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry

(IC)

Immunogen

A synthesized peptide derived from human Kv1.3

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions


Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:50-1:100 Immunocytochemistry (IC): 1:50-1:200

Note: This product is for research use only.

Validation Data




Immunocytochemical analysis of Kv1.3 staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Immunohistochemical analysis of Kv1.3 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-Kv1.3 Rabbit Polyclonal Antibody



Western blotting analysis of Kv1.3 expression in mouse brain (A), rat brain (B) whole cell lysates.