#### **Anti-Cytokeratin 19 Rabbit Polyclonal Antibody**



### **Catalog #: 50451**

#### **Aliases**

Keratin type I cytoskeletal 19; Cytokeratin-19; CK-19; Keratin-19; K19

### **Background**

Gene Name: KRT19 NCBI Gene Entry: 3880 UniProt Entry: P08727

# **Application Information**

Molecular Weight: Predicted, 44 kDa; observed, 44 kDa

Clonality: Rabbit polyclonal antibody Species Reactivity: Human, rat, pig

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry

(IC)

## **Immunogen**

A synthesized peptide derived from human Cytokeratin 19

### **Isotype**

Rabbit IgG

## **Storage Buffer**

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

#### **Storage**


Store at -20 °C for one year.

#### **Recommended Dilutions**


Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:100-1:200 Immunocytochemistry (IC): 1:100-1:500

**Note:** This product is for research use only.


#### Validation Data



Immunocytochemical analysis of Cytokeratin 19 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



Immunohistochemical analysis of Cytokeratin 19 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western blotting analysis of Cytokeratin 19 expression in MCF7 (A), A549 (B), HCC827 (C), HCT116 (D), PC12 (E) whole cell lysates.