#### Anti-S6K1 Rabbit Polyclonal Antibody



## **Catalog #: 50622**

#### **Aliases**

STK14A; Ribosomal protein S6 kinase beta-1; S6K-beta-1; S6K1; 70 kDa ribosomal protein S6 kinase 1; P70S6K1; p70-S6K 1; Ribosomal protein S6 kinase I; Serine/threonine-protein kinase 14A; p70 ribosomal S6 kinase alpha; p70 S6 kinase alpha; p70 S6K-alpha; p70 S6KA

# **Background**

Gene Name: RPS6KB1 NCBI Gene Entry: 6198 UniProt Entry: P23443

# **Application Information**

Molecular Weight: Predicted, 59 kDa; observed, 70,90 kDa

Clonality: Rabbit polyclonal antibody

Species Reactivity: Human, mouse, rat, bovine, chicken, dog, monkey, pig, rabbit

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry

(IC)

## **Immunogen**

A synthesized peptide derived from human S6K1

## **Isotype**

Rabbit IgG

# **Storage Buffer**

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

# **Storage**


Store at -20 °C for one year.

#### **Recommended Dilutions**


Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:100-1:200 Immunocytochemistry (IC): 1:100-1:500

**Note:** This product is for research use only.

#### **Validation Data**




Immunocytochemical analysis of S6K1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Immunohistochemical analysis of S6K1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

# **Anti-S6K1 Rabbit Polyclonal Antibody**



Western blotting analysis of S6K1 expression in MCF7 (A), HeLa (B), HEK293 (C), mouse brain (D) whole cell lysates.