#### **Anti-LRAT Rabbit Polyclonal Antibody**



**Catalog #: 50787** 

#### **Aliases**

Lecithin retinol acyltransferase; Phosphatidylcholine--retinol O-acyltransferase

### **Background**

Gene Name: LRAT NCBI Gene Entry: 9227 UniProt Entry: O95237

# **Application Information**

Molecular Weight: Predicted, 25 kDa; observed, 50 kDa

Clonality: Rabbit polyclonal antibody

Species Reactivity: Human, mouse, rat, monkey, rabbit

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry

(IC)

# **Immunogen**

A synthesized peptide derived from human LRAT

# **Isotype**

Rabbit IgG

# **Storage Buffer**

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

### **Storage**


Store at -20 °C for one year.

### **Recommended Dilutions**


Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:50-1:100 Immunocytochemistry (IC): 1:50-1:200

**Note:** This product is for research use only.

#### Validation Data




Immunocytochemical analysis of LRAT staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Immunohistochemical analysis of LRAT staining in human prostate carcinoma formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

# **Anti-LRAT Rabbit Polyclonal Antibody**



Western blotting analysis of LRAT expression in SHSY5Y (A) whole cell lysates.