Anti-Caspase 9 Rabbit Polyclonal Antibody



Catalog #: 51090

Aliases

MCH6; Caspase-9; CASP-9; Apoptotic protease Mch-6; Apoptotic protease-activating factor 3; APAF-3; ICE-like apoptotic protease 6; ICE-LAP6

Background

Gene Name: CASP9 NCBI Gene Entry: 842 UniProt Entry: P55211

Application Information

Molecular Weight: Predicted, 46 kDa; observed, 47 kDa

Clonality: Rabbit polyclonal antibody

Species Reactivity: Human

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry

(IC)

Immunogen

A synthesized peptide derived from human Caspase 9

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

Storage

Store at -20 °C for one year.


Recommended Dilutions

Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:100-1:200 Immunocytochemistry (IC): 1:100-1:500


Note: This product is for research use only.

Validation Data

TEL: +1-540-855-7041




Immunocytochemical analysis of Caspase 9 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Immunohistochemical analysis of Caspase 9 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-Caspase 9 Rabbit Polyclonal Antibody



Western blotting analysis of Caspase 9 expression in HEK293T (A), Hela (B) whole cell lysates.