Anti-GADD153 Rabbit Polyclonal Antibody



Catalog #: 51107

Aliases

CHOP; CHOP10; GADD153; DNA damage-inducible transcript 3 protein; DDIT-3; C/EBP zeta; C/EBP-homologous protein; CHOP; C/EBP-homologous protein 10; CHOP-10; CCAAT/ enhancer-binding protein homologous protein; Growth arrest and DNA damage-inducible protein GADD153

Background

Gene Name: DDIT3 NCBI Gene Entry: 1649 UniProt Entry: P35638

Application Information

Molecular Weight: Predicted, 19 kDa; observed, 19 kDa

Clonality: Rabbit polyclonal antibody Species Reactivity: Human, monkey, pig

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry

(IC)

Immunogen

A synthesized peptide derived from human GADD153

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.


Storage

Store at -20 °C for one year.


Recommended Dilutions

Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:50-1:100 Immunocytochemistry (IC): 1:50-1:200 **Note:** This product is for research use only.

Validation Data




Immunocytochemical analysis of GADD153 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Immunohistochemical analysis of GADD153 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with

DPX.



Western blotting analysis of GADD153 expression in HCT116 (A), HEK293T (B), A2780 (C), A375 (D) whole cell lysates.