#### **Anti-BMX Rabbit Polyclonal Antibody**



### **Catalog #: 51345**

#### **Aliases**

Cytoplasmic tyrosine-protein kinase BMX; Bone marrow tyrosine kinase gene in chromosome X protein; Epithelial and endothelial tyrosine kinase; ETK; NTK38

## **Background**

Gene Name: BMX NCBI Gene Entry: 660 UniProt Entry: P51813

# **Application Information**

Molecular Weight: Predicted, 78 kDa; observed, 78 kDa

Clonality: Rabbit polyclonal antibody Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry

(IC), immunoprecipitation (IP)

## **Immunogen**

A synthesized peptide derived from human BMX

### **Isotype**

Rabbit IgG

# **Storage Buffer**

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

#### **Storage**


Store at -20 °C for one year.

#### **Recommended Dilutions**


Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:100-1:200 Immunocytochemistry (IC): 1:100-1:500 Immunoprecipitation (IP): 1:10-1:100

**Note:** This product is for research use only.

#### Validation Data




Immunocytochemical analysis of BMX staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



Immunohistochemical analysis of BMX staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

# **Anti-BMX Rabbit Polyclonal Antibody**



Western blotting analysis of BMX expression in mouse brain (A) whole cell lysates.