Anti-UHRF2 Mouse Oligoclonal Antibody



Catalog #: 5156

Aliases

UHRF2; Ubiquitin Like With PHD And Ring Finger Domains 2; RNF107; NIRF; TDRD23; URF2; Ubiquitin-Like With PHD And Ring Finger Domains 2, E3 Ubiquitin Protein Ligase; Ubiquitin-Like-Containing PHD And RING Finger Domains Protein 2; Ubiquitin-Like PHD And RING Finger Domain-Containing Protein 2; RING-Type E3 Ubiquitin Transferase UHRF2; Np95/ICBP90-Like RING Finger Protein; E3 Ubiquitin-Protein Ligase UHRF2; Np95-Like Ring Finger Protein; RING Finger Protein 107; Nuclear Protein 97; MGC33463; Ubiquitin-Like, Containing PHD And RING Finger Domains, 2; Nuclear Zinc Finger Protein NP97; Nuclear Zinc Finger Protein NP97; Np95-Like RING Finger Protein; EC 2.3.2.27; EC 6.3.2

Background

Gene Name: UHRF2

NCBI Gene Entry: 115426 UniProt Entry: Q96PU4

Application Information

Molecular Weight: Predicted, 90 kDa; observed, 90 kDa

Clonality: Mouse oligoclonal antibody Species Reactivity: Human, mouse

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

Immunogen

Recombinant protein of human UHRF2

Isotype

Mouse IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage

Store at -20 °C for one year.


Recommended Dilutions

Western Blotting (WB): 1:500-1:2,500 Flow Cytometry (FCM): 1:2,000


Immunocytochemistry (IC): 1:100-1:1,000

Note: This product is for research use only.


Validation Data



Flow cytometric analysis of UHRF2 expression in HeLa cells using anti-UHRF2 antibody (Cat#5156, 1:2,000). Green, isotype control; red, UHRF2.



Immunocytochemical staining of Hela cells with anti-UHRF2 protein antibody (Cat#5156, 1:1,000). Nuclei were stained blue with DAPI; UHRF2 was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar, 20 μ m.



Western blotting analysis using anti-UHRF2 antibody (Cat#5156). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-UHRF2 antibody (Cat#5156, 1:2,500) and HRP-conjugated goat anti-mouse secondary antibody (Cat#101, 1:20,000) respectively. Image was developed using NaQTM ECL Substrate Kit (Cat#716).