

Anti-DNA Polymerase zeta Rabbit Polyclonal Antibody



Catalog #: 51686

Aliases

POLZ; REV3; DNA polymerase zeta catalytic subunit; Protein reversionless 3-like; REV3-like; hREV3

Background

Gene Name: REV3L

NCBI Gene Entry: [5980](#)

UniProt Entry: [O60673](#)

Application Information

Molecular Weight: Predicted, 352 kDa; observed, 350 kDa

Clonality: Rabbit polyclonal antibody

Species Reactivity: Human, mouse, rat, bovine, chicken, dog, pig

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry (IC)

Immunogen

A synthesized peptide derived from human DNA Polymerase zeta

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions

Western Blotting (WB): 1:500-1:1,000

Immunohistochemistry (IHC): 1:100-1:200

Immunocytochemistry (IC): 1:100-1:500

Note: This product is for research use only.

Validation Data


SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041


ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041


WWW.GENUINBIOTECH.COM



Immunocytochemical analysis of DNA Polymerase zeta staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Immunohistochemical analysis of DNA Polymerase zeta staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western blotting analysis of DNA Polymerase zeta expression in rat lung (A), mouse kidney (B), H1792 (C) whole cell lysates.