Anti-CD61 Rabbit Polyclonal Antibody



Catalog #: 51756

Aliases

GP3A; Integrin beta-3; Platelet membrane glycoprotein IIIa; GPIIIa; CD61

Background

Gene Name: ITGB3 NCBI Gene Entry: 3690 UniProt Entry: P05106

Application Information

Molecular Weight: Predicted, 87 kDa; observed, 87,110 kDa

Clonality: Rabbit polyclonal antibody

Species Reactivity: Human, mouse, rat, monkey

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry

(IC)

Immunogen

A synthesized peptide derived from human CD61

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

Storage


Store at -20 °C for one year.

Recommended Dilutions


Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:50-1:100 Immunocytochemistry (IC): 1:50-1:200

Note: This product is for research use only.


Validation Data



Immunocytochemical analysis of CD61 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).



Immunohistochemical analysis of CD61 staining in human kidney cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western blotting analysis of CD61 expression in mouse liver (A), mouse kidney (B), rat kidney (C), rat testis (D) whole cell lysates.