#### **Anti-FOXO3 Rabbit Polyclonal Antibody**



## **Catalog #: 51868**

#### **Aliases**

FKHRL1; FOXO3A; Forkhead box protein O3; AF6q21 protein; Forkhead in rhabdomyosarcomalike 1

# **Background**

Gene Name: FOXO3 NCBI Gene Entry: 2309 UniProt Entry: O43524

# **Application Information**

Molecular Weight: Predicted, 71 kDa; observed, 97 kDa

Clonality: Rabbit polyclonal antibody Species Reactivity: Human, mouse

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry

(IC)

## **Immunogen**

A synthesized peptide derived from human FOXO3

## **Isotype**

Rabbit IgG

## **Storage Buffer**

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

#### **Storage**


Store at -20 °C for one year.

#### **Recommended Dilutions**


Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:50-1:100 Immunocytochemistry (IC): 1:100-1:500

**Note:** This product is for research use only.

#### Validation Data




Immunocytochemical analysis of FOXO3 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with Alexa Fluor 647-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Immunohistochemical analysis of FOXO3 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

# **Anti-FOXO3 Rabbit Polyclonal Antibody**



Western blotting analysis of FOXO3 expression in PC3 (A), Hela (B), U87MG (C), A549 (D), mouse lung (E) whole cell lysates.