Anti-TNFAIP2 Rabbit Polyclonal Antibody



Catalog #: 52053

Aliases

Tumor necrosis factor alpha-induced protein 2; TNF alpha-induced protein 2; Primary response gene B94 protein

Background

Gene Name: TNFAIP2 NCBI Gene Entry: 7127 UniProt Entry: Q03169

Application Information

Molecular Weight: Predicted, 72 kDa; observed, 72 kDa

Clonality: Rabbit polyclonal antibody

Species Reactivity: Human

Applications Tested: Western blotting (WB), immunohistochemistry (IHC), immunocytochemistry

(IC)

Immunogen

A synthesized peptide derived from human TNFAIP2

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

Storage


Store at -20 °C for one year.

Recommended Dilutions


Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:50-1:200 Immunocytochemistry (IC): 1:50-1:200

Note: This product is for research use only.


Validation Data



Immunocytochemical analysis of TNFAIP2 staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Immunohistochemical analysis of TNFAIP2 staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western blotting analysis of TNFAIP2 expression in A549 (A), A375 (B) whole cell lysates.