

Catalog #: 52792

Aliases

CXCL9; C-X-C Motif Chemokine Ligand 9; SCYB9; Crg-10; Humig; CMK; MIG; Monokine Induced By Gamma Interferon; Monokine Induced By Interferon-Gamma; Gamma-Interferon-Induced Monokine; Chemokine (C-X-C Motif) Ligand 9; Small-Inducible Cytokine B9; C-X-C Motif Chemokine 9; HuMIG

Background

Gene Name: CXCL9

NCBI Gene Entry: [4283](#)

UniProt Entry: [Q07325](#)

Application Information

Molecular Weight: Predicted, 14 kDa; observed, 20 kDa

Clonality: Rabbit polyclonal antibody

Species Reactivity: Mouse

Applications Tested: Western blotting (WB)

Immunogen

A synthesized peptide derived from human CXCL

Isotype

Rabbit IgG

Storage Buffer

Supplied in 0.01M TBS (pH7.4) containing 50% glycerol, 1% BSA and 0.02% Proclin300.

Storage

Store at -20 °C for one year.

Recommended Dilutions

Western Blotting (WB): 1:500-1:2,000

Note: This product is for research use only.

Validation Data


SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM



Western blotting analysis using anti-CXCL9 antibody (Cat#52792). Total lysates were loaded and separated by SDS-PAGE. The blot was incubated with anti-CXCL9 antibody (Cat#52792, 1:1,000) and IRDye800CW-conjugated goat anti-rabbit secondary antibody (1:20,000) respectively.

SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM