Anti-H4C3 Rabbit Monoclonal Antibody



Catalog #: 5318

Aliases

H4C3; H4 Clustered Histone 3; H4/G; DJ221C16.1; HIST1H4C; H4FG; Histone Cluster 1 H4 Family Member C; H4 Histone Family, Member G; Histone Cluster 1, H4c; Histone 1, H4c; Histone H4; H4-16; TEBIVANED1; TEVANED1; HIST1H4A; HIST1H4B; HIST1H4D; HIST1H4E; HIST1H4F; HIST1H4H; HIST1H4I; HIST1H4J; HIST1H4K; HIST1H4L; HIST2H4A; HIST2H4B; HIST2H4; HIST4H4; H4C11; H4C12; H4C13; H4C14; H4C15; H4C16; H4C1; H4C2; H4C4; H4C5; H4C6; H4C8; H4C9; H4/A; H4FA; H4/I; H4FI; H4/B; H4FB; H4/J; H4FJ; H4/C; H4FC; H4/H; H4FH; H4/M; H4FM; H4/E; H4FE; H4/D; H4FD; H4/K; H4FK; H4/N; H4F2; H4FN; H4/O; H4FO

Background

Gene Name: H4C3 NCBI Gene Entry: 8364 UniProt Entry: P62805

Application Information

Molecular Weight: Predicted, 11 kDa; observed, 14 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 25GB920

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM)

Immunogen

A synthesized peptide derived from human Histone H4

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage

Store at -20 °C for one year.


Recommended Dilutions

Western Blotting (WB): 1:1,000-1:5,000


Flow Cytometry (FCM): 1:200-1:2,000

Note: This product is for research use only.

Validation Data



Western blotting analysis using anti-H4C1 antibody (Cat#5318). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-H4C1 antibody (Cat#5318, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using NaQTM ECL Substrate Kit (Cat#716).



Flow cytometric analysis of H4C1 expression in HAP-1 cells using anti-H4C1 antibody (Cat#5318, 1:2,000). Green, isotype control; red, H4C1.

TEL: +1-540-855-7041

SUPPORT@GENUINBIOTECH.COM