#### **Anti-ATIC Mouse Monoclonal Antibody**



### **Catalog #: 5358**

#### **Aliases**

ATIC; 5-Aminoimidazole-4-Carboxamide Ribonucleotide Formyltransferase/IMP Cyclohydrolase; PURH; IMPCHASE; AICARFT; Phosphoribosylaminoimidazolecarboxamide Formyltransferase/IMP Cyclohydrolase; AICAR Transformylase/Inosine Monophosphate Cyclohydrolase; Bifunctional Purine Biosynthesis Protein ATIC; 5-Aminoimidazole-4-Carboxamide-1-Beta-D-Ribonucleotide Transformylase/Inosinicase; AICAR Formyltransferase/IMP Cyclohydrolase Bifunctional Enzyme; Epididymis Secretory Sperm Binding Protein Li 70p; Bifunctional Purine Biosynthesis Protein PURH; AICARFT/IMPCHASE; HEL-S-70p; AICAR

# **Background**

Gene Name: ATIC NCBI Gene Entry: 471 UniProt Entry: P31939

# **Application Information**

Molecular Weight: Predicted, 65 kDa; observed, 65 kDa

Clonality: Mouse monoclonal antibody

Clone ID: 25GB1130

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM)

### Immunogen

Recombinant protein of human ATIC

### **Isotype**

Mouse IgG1

### **Storage Buffer**

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

# **Storage**


Store at -20 °C for one year.

### **Recommended Dilutions**

Western Blotting (WB): 1:400-1:2,000 Flow Cytometry (FCM): 1:2,000


**Note:** This product is for research use only.

#### **Validation Data**



Flow cytometric analysis of ATIC expression in C2C12 cells using anti-ATIC antibody (Cat#5358, 1:2,000). Green, isotype control; red, ATIC.





Western blotting analysis using anti-ATIC antibody (Cat#5358). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-ATIC antibody (Cat#5358, 1:2,000) and HRP-conjugated goat anti-mouse secondary antibody (Cat#101, 1:20,000) respectively. Image was developed using NaQ<sup>TM</sup> ECL Substrate Kit (Cat#716).