

Catalog #: 61360

Aliases

AHR; Aryl Hydrocarbon Receptor; BHLHe76; Class E Basic Helix-Loop-Helix Protein 76; Ah Receptor; BHLHE76; Aromatic Hydrocarbon Receptor; AH-Receptor; RP85; AhR

Background

Gene Name: AHR NCBI Gene Entry: 196 UniProt Entry: P35869

Application Information

Molecular Weight: Predicted, 96 kDa, observed, 105 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 23GB1635

Species Reactivity: Human

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

Immunogen

A synthesized peptide derived from human AhR

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions

Western Blotting (WB): 1:1,000-1:5,000

Flow Cytometry (FCM): 1:2,000


Immunocytochemistry (IC): 1:100-1:1,000

Note: This product is for research use only.

Validation Data


TEL: +1-540-855-7041






Aryl hydrocarbon receptor-Alexa Fluor® 647


Flow cytometric analysis of Aryl hydrocarbon receptor expression in HepG2 cells using Aryl hydrocarbon receptor antibody (Cat#61360, 1:2,000). Green, isotype control; red, Aryl hydrocarbon receptor.



Immunocytochemical staining of HepG2 cells with Aryl hydrocarbon receptor antibody (Cat#61360, 1:1,000). Nuclei were stained blue with DAPI; Aryl hydrocarbon receptor was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 μm.



Western blotting analysis using anti-Aryl hydrocarbon receptor antibody (Cat#61360). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Aryl hydrocarbon receptor antibody (Cat#61360, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQTM ECL Substrate Kit (Cat#226).



Western blotting analysis using anti-Aryl hydrocarbon receptor antibody (Cat#61360). Aryl hydrocarbon receptor expression in wild type (WT) and aryl hydrocarbon receptor shRNA knockdown (KD) Hela cells with 30 μg of total cell lysates. β-Tubulin serves as a loading control. The blot was incubated with anti-Aryl hydrocarbon receptor antibody (Cat#61360, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQTM ECL Substrate Kit (Cat#226).