

Catalog #: 61842

Aliases

TSFM; Ts Translation Elongation Factor, Mitochondrial; Elongation Factor Ts, Mitochondrial; EF-TS; Mitochondrial Elongation Factor Ts; EF-Tsmt; EF-TsMT; EF-TsMt; EFTSMT; EF-Ts; EFTS

Background

Gene Name: TSFM NCBI Gene Entry: 10102 UniProt Entry: P43897

Application Information

Molecular Weight: Predicted, 35 kDa, observed, 32 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 23GB4710

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

Immunogen

A synthesized peptide derived from human TSFM

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions


Western Blotting (WB): 1:1,000-1:5,000

Flow Cytometry (FCM): 1:2,000

Immunocytochemistry (IC): 1:100-1:1,000


Note: This product is for research use only.

Validation Data




Copyright ©2024 Genuin Biotechnologies LLC


Flow cytometric analysis of TSFM expression in HeLa cells using TSFM antibody (Cat#61842, 1:2,000). Green, isotype control; red, TSFM.




Immunocytochemical staining of HeLa cells with TSFM antibody (Cat#61842, 1:1,000). Nuclei were stained blue with DAPI; TSFM was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 µm.




Western blotting analysis using anti-TSFM antibody (Cat#61842). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-TSFM antibody (Cat#61842, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQTM ECL Substrate Kit (Cat#226).



Western blotting analysis using anti-TSFM antibody (Cat#61842). TSFM expression in wild type (WT) and TSFM shRNA knockdown (KD) HeLa cells with 30 μg of total cell lysates. β-Tubulin serves as a loading control. The blot was incubated with anti-TSFM antibody (Cat#61842, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQTM ECL Substrate Kit (Cat#226).



Validation of TSFM knockdown using flow cytometry. Wild-type(WT, Blue) and knockdown(KD, Green) HeLa cells were stained with anti-TSFM antibody (Cat#61842, 1:2,000) and analyzed using BD flow cytometer.



Immunocytochemical staining of HeLa cells using anti-TSFM antibody (Cat#61842, 1:1,000), Top panel: wild-type (WT); Bottom panal: TSFM shRNA knockdown (KD). Nuclei were stained blue with DAPI; TSFM was stained magenta with Alexa Fluor® 647. Scale bar, 20 μ m. Permeabilization: Triton.

TEL: +1-540-855-7041