

**Catalog #: 62313** 

#### **Aliases**

Moesin; Membrane-Organizing Extension Spike Protein; Epididymis Luminal Protein 70; HEL70; IMD50

## **Background**

Gene Name: MSN

NCBI Gene Entry: 4478 UniProt Entry: P26038

## **Application Information**

Molecular Weight: Predicted, 68 kDa, observed, 68 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 24GB265

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

## **Immunogen**

A synthesized peptide derived from human Moesin

## **Isotype**

Rabbit IgG

## **Storage Buffer**

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

## **Storage**

Store at -20 °C for one year.

### **Recommended Dilutions**

Western Blotting (WB): 1:1,000-1:5,000


Flow Cytometry (FCM): 1:2,000

Immunocytochemistry (IC): 1:100-1:1,000


**Note:** This product is for research use only.

#### Validation Data

TEL: +1-540-855-7041




Flow cytometric analysis of Moesin expression in HepG2 cells using Moesin antibody (Cat#62313, 1:2,000). Green, isotype control; red, Moesin.




Immunocytochemical staining of Hela cells with anti-Moesin antibody (Cat#62313, 1:1,000). Nuclei were stained blue with DAPI; Moesin was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain:

Medium. Scale bar: 20 µm.




Western blotting analysis using anti-Moesin antibody (Cat#62313). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Moesin antibody (Cat#62313, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQ<sup>TM</sup> ECL Substrate Kit (Cat#226).




Copyright @2024 Genuin Biotechnologies LLC

Western blotting analysis using anti-Moesin antibody (Cat#62313). Moesin expression in wild type (WT) and Moesin shRNA knockdown (KD) HeLa cells with 20 μg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with anti-Moesin antibody (Cat#62313, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using NaQ<sup>TM</sup> ECL Substrate Kit (Cat#716).



Validation of Moesin knockdown using flow cytometry. Wild-type(WT, Blue) and knockdown(KD, Green) HeLa cells were stained with anti-Moesin antibody (Cat#62313, 1:2,000) and analyzed using BD flow cytometer.



Immunocytochemical staining of HeLa cells using anti-Moesin antibody (Cat#62313, 1:1,000), Top panel: wild-type (WT); Bottom panal: Moesin shRNA knockdown (KD). Nuclei were stained blue with DAPI; Moesin was stained magenta with Alexa Fluor® 647. Scale bar, 20 μm.