

Catalog #: 68610

Aliases

SRC; SRC Proto-Oncogene, Non-Receptor Tyrosine Kinase; C-Src; SRC1; ASV; V-Src Avian Sarcoma (Schmidt-Ruppin A-2) Viral Oncogene Homolog; Proto-Oncogene Tyrosine-Protein Kinase Src; Proto-Oncogene C-Src; EC 2.7.10.2; P60-Src; Protooncogene SRC, Rous Sarcoma; Tyrosine-Protein Kinase SRC-1; Tyrosine Kinase Pp60c-Src; Pp60c-Src; EC 2.7.10; THC6

Background

Gene Name: SRC

NCBI Gene Entry: 6714 UniProt Entry: P12931

Application Information

Molecular Weight: Predicted, 60 kDa, observed, 60 kDa

Clonality: Rabbit monoclonal antibody

Clone ID: 23GB1110

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), flow cytometry (FCM), immunocytochemistry (IC)

Immunogen

A synthesized peptide derived from human Src

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions


Western Blotting (WB): 1:1,000-1:5,000

Flow Cytometry (FCM): 1:2,000


Immunocytochemistry (IC): 1:100-1:1,000

Note: This product is for research use only.

Validation Data




Flow cytometric analysis of Src expression in C2C12 cells using Src antibody (Cat#68610, 1:2,000). Green, isotype control; red, Src.




Immunocytochemical staining of HeLa cells with Src antibody (Cat#68610, 1:1,000). Nuclei were stained blue with DAPI; Src was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar: 20 μm.

TEL: +1-540-855-7041



Western blotting analysis using anti-Src antibody (Cat#68610). Total cell lysates (30 μg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-Src antibody (Cat#68610, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:50,000) respectively. Image was developed using FeQTM ECL Substrate Kit (Cat#226).



Western blotting analysis using anti-Src antibody (Cat#68610). Src expression in wild type (WT) and Src shRNA knockdown (KD) HT-1080 cells with 30 μg of total cell lysates. β-Tubulin serves as a loading control. The blot was incubated with anti-Src antibody (Cat#68610, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000) respectively. Image was developed using FeQTM ECL Substrate Kit (Cat#226).