# **Human DHFR Knockdown Cell Line (WB-Validated)**



**Catalog #: C61398** 

#### **Aliases**

DHFR; Dihydrofolate Reductase; DHFR1; EC 1.5.1.3; DHFRP1; DYR

## **Background**

Gene Name: DHFR NCBI Gene Entry: 1719

## **Storage**

Store at liquid nitrogen for 1 year.

## **Kit Components**

- 1. Human DHFR Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

#### **Parental Cell Line**

Human cell line supplied by the client

### **Validation Methods**

RT-qPCR, Western blotting (WB)

### **Shipping**

Shipped on Dry Ice.


### **Instructions For Use**

This knockdown cell line should be paired with wild-type cell line for use.

**Note:** This product is for research use only.


#### **Validation Data**

# **Human DHFR Knockdown Cell Line (WB-Validated)**



Genotype	Ct Value
Wild-Type	19.51
Knock-Down	23.02
ΔCt (Ct <sub>KD</sub> -Ct <sub>WT</sub> )	3.51 80 And the state of the st
% mRNA Reduction	<b>J</b> 91%

RT-qPCR analysis. HeLa cells were infected with DHFR-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers.  $\Delta$ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula:  $(1-1/2\Delta$ Ct) x 100%.



Western blotting analysis. DHFR protein expression in wild-type (WT) and shRNA knockdown

#### PAGE 3

# **Human DHFR Knockdown Cell Line (WB-Validated)**

(KD) HeLa cells was detected using Western blotting.  $\beta$ -Tubulin served as a loading control. The blots were incubated with primary antibodies (Cat#61398, 1:5,000) against DHFR and  $\beta$ -Tubulin, respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000). Images were developed using FeQ<sup>TM</sup> ECL Substrate Kit (Cat#226).