Human HDAC8 Knockdown Cell Line (WB-Validated)



Catalog #: C61423

Aliases

HDAC8; Histone Deacetylase 8; HDACL1; KDAC8; RPD3; Protein Decrotonylase HDAC8; Histone Deacetylase-Like 1; Protein Deacetylase HDAC8; MRXS6; HD8; WTS; Wilson-Turner X-Linked Mental Retardation Syndrome; EC 3.5.1.98; EC 3.5.1.-; CDA07; CDLS5

Background

Gene Name: HDAC8 NCBI Gene Entry: 55869

Storage

Store at liquid nitrogen for 1 year.

Kit Components

- 1. Human HDAC8 Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

Parental Cell Line

Human cell line supplied by the client

Validation Methods

RT-qPCR, Western blotting (WB)

Shipping

Shipped on Dry Ice.


Instructions For Use

This knockdown cell line should be paired with wild-type cell line for use.

Note: This product is for research use only.


Validation Data

Human HDAC8 Knockdown Cell Line (WB-Validated)



Genotype	Ct Value
Wild-Type	21.22
Knock-Down	22.47
$\Delta Ct (Ct_{KD}-Ct_{WT})$	1.25
% mRNA Reduction	↓ 58%

RT-qPCR analysis. HeLa cells were infected with HDAC8-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers. Δ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula: $(1-1/2\Delta$ Ct) x 100%.



Western blotting analysis.HDAC8 protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting. Hsp90 α served as a loading control. The blots were incubated with primary antibodies against HDAC8 and Hsp90 α, respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody. Images were developed using NaQTM ECL Substrate Kit.