# **Human ENG Knockdown Cell Line (WB-Validated)**



**Catalog #: C61450** 

#### **Aliases**

ENG; Endoglin; END; HHT1; CD105 Antigen; CD105; ORW1; Osler-Rendu-Weber Syndrome 1; Soluble Endoglin; ORW

# **Background**

Gene Name: ENG

NCBI Gene Entry: 2022

## **Storage**

Store at liquid nitrogen for 1 year.

## **Kit Components**

- 1. Human ENG Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

#### **Parental Cell Line**

Human cell line supplied by the client

### **Validation Methods**

RT-qPCR, Western blotting (WB)

# **Shipping**

Shipped on Dry Ice.


### **Instructions For Use**

This knockdown cell line should be paired with wild-type cell line for use.

**Note:** This product is for research use only.


#### **Validation Data**

# **Human ENG Knockdown Cell Line (WB-Validated)**



Genotype	Ct Value
Wild-Type	21.50
Knock-Down	26.29
$\Delta Ct (Ct_{KD}-Ct_{WT})$	4.79
% mRNA Reduction	<b>♣ 96%</b>

RT-qPCR analysis. HeLa cells were infected with ENG-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers.  $\Delta$ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula:  $(1-1/2\Delta$ Ct) x 100%.



Western blotting analysis. ENG protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting. β-Tubulin served as a loading control. The blots were incubated with primary antibodies (Cat#61450, 1:5,000) against ENG and β-Tubulin, respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000). Images were developed using FeQ<sup>TM</sup> ECL Substrate Kit (Cat#226).