# **Human RAF1 Knockdown Cell Line (WB-Validated)**



**Catalog #: C61512** 

#### **Aliases**

RAF1; Raf-1 Proto-Oncogene, Serine/Threonine Kinase; Raf-1; CRAF; RAF Proto-Oncogene Serine/Threonine-Protein Kinase; V-Raf-1 Murine Leukemia Viral Oncogene Homolog 1; C-Raf Proto-Oncogene, Serine/Threonine Kinase; Proto-Oncogene C-RAF; EC 2.7.11.1; C-Raf; V-Raf-1 Murine Leukemia Viral Oncogene-Like Protein 1; Raf Proto-Oncogene Serine/Threonine Protein Kinase; Oncogene RAF1; EC 2.7.11; CMD1NN; C-RAF; RAF-1; CRaf; NS5; RAF

## **Background**

Gene Name: RAF1 NCBI Gene Entry: 5894

# **Storage**

Store at liquid nitrogen for 1 year.

# **Kit Components**

- 1. Human RAF1 Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

### **Parental Cell Line**

Human cell line supplied by the client

### **Validation Methods**

RT-qPCR, Western blotting (WB)

## **Shipping**

Shipped on Dry Ice.


### **Instructions For Use**

This knockdown cell line should be paired with wild-type cell line for use.

**Note:** This product is for research use only.


#### Validation Data

# **Human RAF1 Knockdown Cell Line (WB-Validated)**



Genotype	Ct Value
Wild-Type	19.27
Knock-Down	20.49
$\Delta$ Ct (Ct <sub>KD</sub> -Ct <sub>WT</sub> )	1.22
% mRNA Reduction	<b>↓</b> 57%

RT-qPCR analysis. HeLa cells were infected with RAF1-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers.  $\Delta$ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula:  $(1-1/2\Delta$ Ct) x 100%.



#### PAGE 3

# **Human RAF1 Knockdown Cell Line (WB-Validated)**

Western blotting analysis. RAF1 protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting.  $\beta$ -Tubulin served as a loading control. The blots were incubated with primary antibodies (Cat#61510, 1:5,000) against RAF1 and  $\beta$ -Tubulin , respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody (Cat#201, 1:20,000). Images were developed using FeQ<sup>TM</sup> ECL Substrate Kit (Cat#226).