# **Human DNM2 Knockdown Cell Line (WB-Validated)**



**Catalog #: C61929** 

#### **Aliases**

DNM2; Dynamin 2; DYN2; Dynamin II; DI-CMTB; CMTDIB; CMTDI1; DYNII; CMT2M; EC 3.6.5.5; Dynamin-2; Cytoskeletal Protein; LCCS5

## **Background**

Gene Name: DNM2 NCBI Gene Entry: 1785

## **Storage**

Store at liquid nitrogen for 1 year.

## Kit Components

- 1. Human DNM2 Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

### **Parental Cell Line**

Human cell line supplied by the client

### Validation Methods

RT-qPCR, Western blotting (WB)

# **Shipping**

Shipped on Dry Ice.


## **Instructions For Use**

This knockdown cell line should be paired with wild-type cell line for use.

**Note:** This product is for research use only.


#### **Validation Data**

# **Human DNM2 Knockdown Cell Line (WB-Validated)**



Genotype	Ct Value
Wild-Type	20.20
Knock-Down	21.77
$\Delta Ct (Ct_{KD}-Ct_{WT})$	1.57
% mRNA Reduction	<b>4</b> 66%

RT-qPCR analysis. HeLa cells were infected with DNM2-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers.  $\Delta$ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula:  $(1-1/2\Delta$ Ct) x 100%.



Western blotting analysis. DNM2 protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting.  $\beta$ -Tubulin served as a loading control. The blots were incubated with primary antibodies against DNM2 and  $\beta$ -Tubulin, respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody. Images were developed using FeQ<sup>TM</sup> ECL Substrate Kit.