# **Human DYNLL1 Knockdown Cell Line (WB-Validated)**



**Catalog #: C61946** 

#### **Aliases**

DYNLL1; Dynein Light Chain LC8-Type 1; DLC1; DLC8; PIN; Hdlc1; DNCL1; LC8; Protein Inhibitor Of Neuronal Nitric Oxide Synthase; Dynein, Cytoplasmic, Light Polypeptide 1; Dynein Light Chain 1, Cytoplasmic; 8 KDa Dynein Light Chain; DNCLC1; Cytoplasmic Dynein Light Polypeptide; HDLC1; LC8a

## **Background**

Gene Name: DYNLL1 NCBI Gene Entry: 8655

## **Storage**

Store at liquid nitrogen for 1 year.

## **Kit Components**

- 1. Human DYNLL1 Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

### **Parental Cell Line**

Human cell line supplied by the client

### **Validation Methods**

RT-qPCR, Western blotting (WB)

## **Shipping**

Shipped on Dry Ice.


### **Instructions For Use**

This knockdown cell line should be paired with wild-type cell line for use.


**Note:** This product is for research use only.

#### Validation Data

# **Human DYNLL1 Knockdown Cell Line (WB-Validated)**



RT-qPCR analysis. HeLa cells were infected with DYNLL1-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers.  $\Delta$ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula:  $(1-1/2\Delta$ Ct) x 100%.



Western blotting analysis. DYNLL1 protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting.  $\beta$ -Tubulin served as a loading control. The blots were incubated with primary antibodies against DYNLL1 and  $\beta$ -Tubulin, respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody. Images were developed using FeQ<sup>TM</sup> ECL Substrate Kit.