# **Human NLK Knockdown Cell Line (WB-Validated)**



**Catalog #: C62384** 

#### **Aliases**

NLK; Nemo Like Kinase; Serine/Threonine-Protein Kinase NLK; Nemo-Like Kinase; EC 2.7.11.24; Protein LAK1; EC 2.7.11; LAK1

## **Background**

Gene Name: NLK

NCBI Gene Entry: 51701

## **Storage**

Store at liquid nitrogen for 1 year.

## **Kit Components**

- 1. Human NLK Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

### **Parental Cell Line**

Human cell line supplied by the client

### Validation Methods

RT-qPCR, Western blotting (WB)

# **Shipping**

Shipped on Dry Ice.


## **Instructions For Use**

This knockdown cell line should be paired with wild-type cell line for use.

**Note:** This product is for research use only.


#### **Validation Data**

# **Human NLK Knockdown Cell Line (WB-Validated)**



Genotype	Ct Value ্র
Wild-Type	20.89
Knock-Down	21.98
ΔCt (Ct <sub>KD</sub> -Ct <sub>WT</sub> )	1.09
% mRNA Reduction	<b>↓</b> 53%

RT-qPCR analysis. HT-1080 cells were infected with NLK-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers.  $\Delta$ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula:  $(1-1/2\Delta$ Ct) x 100%.



Western blotting analysis. NLK protein expression in wild-type (WT) and shRNA knockdown (KD) HT-1080 cells was detected using Western blotting. Hsp90  $\alpha$  served as a loading control. The blots were incubated with primary antibodies against NLK and Hsp90  $\alpha$ , respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody. Images were developed using FeQ<sup>TM</sup> ECL Substrate Kit.