

Human PPIA Knockdown Cell Line (WB-Validated)



Catalog #: C62452

Aliases

PPIA; Peptidylprolyl Isomerase A; CYPA; Peptidyl-Prolyl Cis-Trans Isomerase A; Cyclosporin A-Binding Protein; Cyclophilin A; Rotamase A; EC 5.2.1.8; PPIase A; Epididymis Secretory Sperm Binding Protein Li 69p; Peptidylprolyl Isomerase A (Cyclophilin A); T Cell Cyclophilin; HEL-S-69p; CYPH

Background

Gene Name: PPIA

NCBI Gene Entry: [5478](#)

Storage

Store at liquid nitrogen for 1 year.

Kit Components

1. Human PPIA Knockdown Cell Line (Wb-Validated)
2. Wild-type cell line

Parental Cell Line

Human cell line supplied by the client

Validation Methods

RT-qPCR, Western blotting (WB)

Shipping

Shipped on Dry Ice.

Instructions For Use

This knockdown cell line should be paired with wild-type cell line for use.

Note: This product is for research use only.

Validation Data


SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS


SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM



Genotype	Ct Value
Wild-Type	19.26
Knock-Down	21.13
$\Delta Ct (Ct_{KD}-Ct_{WT})$	1.87
% mRNA Reduction	↓ 73%

RT-qPCR analysis. HeLa cells were infected with PPIA-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers. $\Delta Ct (Ct_{KD}-Ct_{WT})$ was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula: $(1-1/2^{\Delta Ct}) \times 100\%$.



Western blotting analysis. PPIA protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting. β -Tubulin served as a loading control. The blots were incubated with primary antibodies against PPIA and β -Tubulin, respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody. Images were developed using FeQ™ ECL Substrate Kit.