Human TACC3 Knockdown Cell Line (WB-Validated)

Catalog #: C62642

Aliases

TACC3; Transforming Acidic Coiled-Coil Containing Protein 3; ERIC-1; ERIC1; Maskin; Tacc4; Transforming Acidic Coiled-Coil-Containing Protein 3; Transforming, Acidic Coiled-Coil Containing Protein 3

Background

Gene Name: TACC3
NCBI Gene Entry: 10460

Storage

Store at liquid nitrogen for 1 year.

Kit Components

1. Human TACC3 Knockdown Cell Line (Wb-Validated)
2. Wild-type cell line

Parental Cell Line

Human cell line supplied by the client

Validation Methods

RT-qPCR, Western blotting (WB)

Shipping

Shipped on Dry Ice.

Instructions For Use

This knockdown cell line should be paired with wild-type cell line for use.

Note: This product is for research use only.

Validation Data

Human TACC3 Knockdown Cell Line (WB-Validated)



Genotype	Ct Value
Wild-Type	21.51
Knock-Down	24.75
ΔCt (Ct_{KD}-Ct_{WT})	3.24
% mRNA Reduction	⬇ 89%

RT-qPCR analysis. HeLa cells were infected with TACC3-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers. ΔCt (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula: (1-1/2ΔCt) x 100%.



Western blotting analysis. TACC3 protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting. β-Tubulin served as a loading control. The blots were incubated with primary antibodies against TACC3 and β-Tubulin, respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody. Images were developed using FeQ™ ECL Substrate Kit.