Human TNFAIP8 Knockdown Cell Line (WB-Validated)



Catalog #: C62659

Aliases

TNFAIP8; TNF Alpha Induced Protein 8; MDC-3.13; SCC-S2; GG2-1; Head And Neck Tumor And Metastasis-Related Protein; Tumor Necrosis Factor Alpha-Induced Protein 8; NF-Kappa-B-Inducible DED-Containing Protein; TNF-Induced Protein GG2-1; NDED; Tumor Necrosis Factor, Alpha-Induced Protein 8; Tumor Necrosis Factor, Alpha Induced Protein 8; TNF Alpha-Induced Protein 8; SCCS2

Background

Gene Name: TNFAIP8 NCBI Gene Entry: 25816

Storage

Store at liquid nitrogen for 1 year.

Kit Components

- 1. Human TNFAIP8 Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

Parental Cell Line

Human cell line supplied by the client

Validation Methods

RT-qPCR, Western blotting (WB)

Shipping

Shipped on Dry Ice.


Instructions For Use

This knockdown cell line should be paired with wild-type cell line for use.

Note: This product is for research use only.


Validation Data

Human TNFAIP8 Knockdown Cell Line (WB-Validated)



Genotype	Ct Value
Wild-Type	24.40
Knock-Down	26.33
$\Delta Ct (Ct_{KD}-Ct_{WT})$	1.93
% mRNA Reduction	↓ 74%

RT-qPCR analysis. HeLa cells were infected with TNFAIP8-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers. Δ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula: $(1-1/2\Delta$ Ct) x 100%.



Western blotting analysis. TNFAIP8 protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting. β -Tubulin served as a loading control. The blots were incubated with primary antibodies against TNFAIP8 and β -Tubulin, respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody. Images were developed using FeQTM ECL Substrate Kit.