# **Human FGF2 Knockdown Cell Line (WB-Validated)**



**Catalog #: C63186** 

#### **Aliases**

FGF2; Fibroblast Growth Factor 2; FGFB; Fibroblast Growth Factor 2 (Basic); Heparin-Binding Growth Factor 2; HBGF-2; FGF-2; BFGF; Basic Fibroblast Growth Factor BFGF; Basic Fibroblast Growth Factor; Prostatropin

## **Background**

Gene Name: FGF2 NCBI Gene Entry: 2247

## **Storage**

Store at liquid nitrogen for 1 year.

## **Kit Components**

- 1. Human FGF2 Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

### **Parental Cell Line**

Human cell line supplied by the client

### Validation Methods

RT-qPCR, Western blotting (WB)

# **Shipping**

Shipped on Dry Ice.


## **Instructions For Use**

This knockdown cell line should be paired with wild-type cell line for use.

**Note:** This product is for research use only.


#### **Validation Data**

# **Human FGF2 Knockdown Cell Line (WB-Validated)**



Genotype	Ct Value
Wild-Type	26.37 sign
Knock-Down	<b>29.71</b>
∆Ct (CtKD-CtWT)	3.34
% mRNA	opyright ©
Reduction	90% <sup>§</sup>

RT-qPCR analysis. HeLa cells were infected with FGF2-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers.  $\Delta$ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula:  $(1-1/2\Delta$ Ct) x 100%.



Western blotting analysis. FGF2 protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting.  $\beta$ -Tubulin served as a loading control. The blots were incubated with primary antibodies against FGF2 and  $\beta$ -Tubulin, respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody. Images were developed using FeQ<sup>TM</sup> ECL Substrate Kit.