# **Human GRN Knockdown Cell Line (WB-Validated)**



**Catalog #: C63575** 

#### **Aliases**

GRN; Granulin Precursor; PCDGF; PGRN; Progranulin; CLN11; PC Cell-Derived Growth Factor; Glycoprotein Of 88 Kda; Glycoprotein 88; Proepithelin; Acrogranin; GP88; PEPI; Epithelin Precursor; Granulin-Epithelin; Epithelin; Granulins; GEP; FTD2

## **Background**

Gene Name: GRN

NCBI Gene Entry: 2896

### **Storage**

Store at liquid nitrogen for 1 year.

## **Kit Components**

- 1. Human GRN Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

#### **Parental Cell Line**

Human cell line supplied by the client

### Validation Methods

RT-qPCR, Western blotting (WB)

## **Shipping**

Shipped on Dry Ice.


## **Instructions For Use**

This knockdown cell line should be paired with wild-type cell line for use.


**Note:** This product is for research use only.

#### Validation Data

## **Human GRN Knockdown Cell Line (WB-Validated)**



RT-qPCR analysis. HeLa cells were infected with GRN-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers.  $\Delta$ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula:  $(1-1/2\Delta$ Ct) x 100%.



Western blotting analysis. GRN protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting. Hsp90  $\alpha$  served as a loading control. The blots were incubated with primary antibodies against GRN and Hsp90  $\alpha$ , respectively, followed by incubating with HRP-conjugated goat anti-mouse secondary antibody. Images were developed using FeQ<sup>TM</sup> ECL Substrate Kit.