

Human BCAT1 Knockdown Cell Line (WB-Validated)



Catalog #: C63802

Aliases

BCAT1; Branched Chain Amino Acid Transaminase 1; BCATc; BCT1; Branched-Chain-Amino-Acid Aminotransferase, Cytosolic; Branched Chain Aminotransferase 1, Cytosolic; EC 2.6.1.42; ECA39; Branched Chain Amino-Acid Transaminase 1, Cytosolic; Placental Protein 18; Protein ECA39; PNAS121; BCAT(C); MECA39; PP18

Background

Gene Name: BCAT1

NCBI Gene Entry: [586](#)

Storage

Store at liquid nitrogen for 1 year.

Kit Components

1. Human BCAT1 Knockdown Cell Line (Wb-Validated)
2. Wild-type cell line

Parental Cell Line

Human cell line supplied by the client

Validation Methods

RT-qPCR, Western blotting (WB)

Shipping

Shipped on Dry Ice.

Instructions For Use

This knockdown cell line should be paired with wild-type cell line for use.

Note: This product is for research use only.

Validation Data


SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041


ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM



RT-qPCR analysis. HeLa cells were infected with BCAT1-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers. $\Delta Ct (Ct_{KD}-Ct_{WT})$ was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula: $(1-1/2^{\Delta Ct}) \times 100\%$.



Western blotting analysis. BCAT1 protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting. Hsp90 α served as a loading control. The blots were incubated with primary antibodies against BCAT1 and Hsp90 α, respectively, followed by incubating with HRP-conjugated goat anti-mouse secondary antibody. Images were developed

Human BCAT1 Knockdown Cell Line (WB-Validated)

PAGE 3

using FeQ™ ECL Substrate Kit.

SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM