# **Human UCHL5 Knockdown Cell Line (WB-Validated)**



**Catalog #: C65519** 

#### **Aliases**

UCHL5; Ubiquitin C-Terminal Hydrolase L5; UCH37; CGI-70; INO80R; Ubiquitin Carboxyl-Terminal Hydrolase Isozyme L5; Ubiquitin Carboxyl-Terminal Hydrolase L5; Ubiquitin C-Terminal Hydrolase UCH37; Ubiquitin Thioesterase L5; INO80 Complex Subunit R; EC 3.4.19.12; UCH-L5; Ubiquitin Carboxyl-Terminal Esterase L5

## **Background**

Gene Name: UCHL5 NCBI Gene Entry: 51377

## **Storage**

Store at liquid nitrogen for 1 year.

# **Kit Components**

- 1. Human UCHL5 Knockdown Cell Line (Wb-Validated)
- 2. Wild-type cell line

### **Parental Cell Line**

Human cell line supplied by the client

### **Validation Methods**

RT-qPCR, Western blotting (WB)

## **Shipping**

Shipped on Dry Ice.


### **Instructions For Use**

This knockdown cell line should be paired with wild-type cell line for use.


**Note:** This product is for research use only.

#### Validation Data

# **Human UCHL5 Knockdown Cell Line (WB-Validated)**



RT-qPCR analysis. HeLa cells were infected with UCHL5-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers.  $\Delta$ Ct (CtKD-CtWT) was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula:  $(1-1/2\Delta$ Ct) x 100%.



Western blotting analysis.UCHL5 protein expression in wild-type (WT) and shRNA knockdown (KD) HeLa cells was detected using Western blotting. Hsp90  $\alpha$  served as a loading control. The blots were incubated with primary antibodies against UCHL5 and Hsp90  $\alpha$ , respectively, followed by incubating with HRP-conjugated goat anti-rabbit secondary antibody. Images were developed

TEL: +1-540-855-7041

#### PAGE 3

# **Human UCHL5 Knockdown Cell Line (WB-Validated)**

using NaQ<sup>TM</sup> ECL Substrate Kit.