#### **Anti-His Tag Mouse Monoclonal Antibody**



**Catalog #: T008** 

#### **Aliases**

6 His epitope tag; Hexa His tag; HHHHHHH epitope tag; HHHHHHH tag; His tag

## **Background**

The His-Tag (or Polyhistidine tag) is a short amino acid sequence, typically composed of six to ten consecutive histidine residues (e.g., His<sub>6</sub>), genetically engineered onto either the N-or C-terminus of a recombinant protein. Its primary function is to facilitate the purification of the tagged protein using immobilized metal affinity chromatography (IMAC). Histidine residues possess an imidazole side chain that exhibits high affinity for certain divalent metal ions, such as nickel (Ni<sup>2+</sup>), cobalt (Co<sup>2+</sup>), copper (Cu<sup>2+</sup>), or zinc (Zn<sup>2+</sup>). This specific interaction allows the His-tagged protein to bind tightly to a chromatography resin charged with one of these metal ions.

# **Application Information**

Molecular Weight: Recombinant protein dependent

Clonality: Mouse monoclonal antibody

Clone ID: 25GB7085

Species Reactivity: Recombinant protein Applications Tested: Western blotting (WB)

#### Immunogen

Recombinant protein of His tag

## **Isotype**

Mouse IgG1

### **Storage Buffer**

Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.

# **Storage**


Store at -20 °C for one year.

### **Recommended Dilutions**

Western Blotting (WB): 1:1,000-1:5,000

Note: This product is for research use only.

#### **Validation Data**



Western blotting analysis using anti-His-tag antibody (Cat#T008). HHHHHH tagged-fusion protein (0.50  $\mu$ g) expressed by E. coli were loaded and separated by SDS-PAGE. The blot was incubated with anti-His-tag antibody (Cat#T008, 1:5,000) and HRP-conjugated goat anti-mouse secondary antibody (Cat#101, 1:20,000) respectively. Image was developed using FeQ<sup>TM</sup> ECL Substrate Kit (Cat#226).