

Anti-Phospho-Histone Deacetylase 7 (S155) Rabbit Polyclonal Antibody



Catalog #: U0636

Aliases

HDAC7A; Histone deacetylase 7; HD7; Histone deacetylase 7A; HD7a

Background

Gene Name: HDAC7

NCBI Gene Entry: [51564](#)

UniProt Entry: [Q8WUI4](#)

Application Information

Molecular Weight: Predicted, 102 kDa; observed, 105 kDa

Clonality: Rabbit polyclonal antibody

Species Reactivity: Human, mouse, rat, zebrafish

Applications Tested: Western blotting (WB), immunocytochemistry (IC)

Immunogen

A synthesized peptide derived from human Phospho-Histone Deacetylase 7 (S155)

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions

Western Blotting (WB): 1:500-1:1,000

Immunocytochemistry (IC): 1:100-1:500

Note: This product is for research use only.

Validation Data

SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041


ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041


WWW.GENUINBIOTECH.COM

Anti-Phospho-Histone Deacetylase 7 (S155) Rabbit Polyclonal Antibody

PAGE 2



Immunocytochemical analysis of Histone Deacetylase 7 (Phospho-S155) staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



Western blotting analysis of Histone Deacetylase 7 (Phospho-S155) expression in HEK293T (A), HeLa (B), U2OS (C), rat lung (D) whole cell lysates.

SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM