# **Anti-Phospho-CDK7 (T170) Rabbit Polyclonal Antibody**



#### **Catalog #: U0795**

#### **Aliases**

CAK; CAK1; CDKN7; MO15; STK1; Cyclin-dependent kinase 7; 39 kDa protein kinase; p39 Mo15; CDK-activating kinase 1; Cell division protein kinase 7; Serine/threonine-protein kinase 1; TFIIH basal transcription factor complex kinase subunit

### **Background**

Gene Name: CDK7 NCBI Gene Entry: 1022 UniProt Entry: P50613

## **Application Information**

Molecular Weight: Predicted, 39 kDa; observed, 40 kDa

Clonality: Rabbit polyclonal antibody Species Reactivity: Human, mouse

Applications Tested: Western blotting (WB), immunohistochemistry (IHC)

#### **Immunogen**

A synthesized peptide derived from human Phospho-CDK7 (T170)

### **Isotype**

Rabbit IgG

#### **Storage Buffer**

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

#### **Storage**

Store at -20 °C for one year.


#### **Recommended Dilutions**

Western Blotting (WB): 1:500-1:1,000 Immunohistochemistry (IHC): 1:50-1:100


**Note:** This product is for research use only.

#### **Validation Data**

## **Anti-Phospho-CDK7 (T170) Rabbit Polyclonal Antibody**



Immunohistochemical analysis of CDK7 (Phospho-T170) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western blotting analysis of CDK7 (Phospho-T170) expression in Hela (A), HEK293T (B), A549 (C), MCF7 (D), Raw264.7 (E), mouse spleen (F) whole cell lysates.