

Anti-Phospho-Histone H4 (S80) Rabbit Polyclonal Antibody



Catalog #: U1039

Aliases

H4/A; H4FA; H4/I; H4FI; H4/G; H4FG; H4/B; H4FB; H4/J; H4FJ; H4/C; H4FC; H4/H; H4FH; H4/M; H4FM; H4/E; H4FE; H4/D; H4FD; H4/K; H4FK; H4/N; H4F2; H4FN; HIST2H4; H4/O; H4FO; Histone H4

Background

Gene Name: HIST1H3A/3B/3C/3D/3E/3F/3G/3H/3I/3J/3A/3C/3D/H3F3A/H3F3B

NCBI Gene Entry: [121504](#)

UniProt Entry: [P62805](#)

Application Information

Molecular Weight: Predicted, 11 kDa; observed, 14 kDa

Clonality: Rabbit polyclonal antibody

Species Reactivity: Human, mouse, rat

Applications Tested: Western blotting (WB), immunohistochemistry (IHC)

Immunogen

A synthesized peptide derived from human Phospho-Histone H4 (S80)

Isotype

Rabbit IgG

Storage Buffer

Supplied in PBS (pH 7.3) containing 30% glycerol, and 0.01% sodium azide.

Storage

Store at -20 °C for one year.

Recommended Dilutions

Western Blotting (WB): 1:1,000-1:2,000

Immunohistochemistry (IHC): 1:100-1:200

Note: This product is for research use only.

Validation Data


SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041


ORDERS

SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM



Immunohistochemical analysis of Histone H4 (Phospho-S80) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western blotting analysis of Histone H4 (Phospho-S80) expression in HeLa (A) whole cell lysates.