

WB-Validated CDC16 Lentiviral shRNA Knockdown Kit



Catalog #: V61386

Aliases

CDC16; Cell Division Cycle 16; ANAPC6; APC6; CUT9; Cell Division Cycle Protein 16 Homolog; Anaphase-Promoting Complex, Subunit 6; Cyclosome Subunit 6; CDC16Hs; CDC16 (Cell Division Cycle 16, *S. Cerevisiae*, Homolog); CDC16 Cell Division Cycle 16 Homolog (*S. Cerevisiae*); Cell Division Cycle 16 Homolog (*S. Cerevisiae*); Anaphase-Promoting Complex Subunit 6; Cell Division Cycle 16 Homolog; CDC16 Homolog

Background

Gene Name: CDC16
NCBI Gene Entry: [8881](#)

Storage

Store at -80 °C for one year.

Kit Components

1. WB-Validated CDC16 shRNA lentiviral particles (4 mL)
- 2 . Non-Target shRNA lentiviral particles (4 mL)
3. Verification Tool: KD-Validated Anti-CDC16 Rabbit mAb #61386 (5 µL)

Tested Cell Line

HeLa

Validation Methods

RT-qPCR; Western Blotting (WB)

Shipping

Shipped with dry ice. Immediately store the product in a standard freezer at -80°C upon receipt.

Instructions For Use

The following protocol uses HeLa cell as an example assuming your cell culture medium is DMEM-based.

1. Release 0.5 million HeLa cells into a 35 mm tissue culture dish in 2 mL of the growth medium (DMEM containing 10% FBS and 1% pen/strep). Cell density should reach 50-60% confluence the following day.
2. 24 h after cell release, pre-warm the shRNA lentiviral medium to 37°C.
3. Discard 1 mL of the original growth medium of the 35 mm dish.

SUPPORT

SUPPORT@GENUINBIOTECH.COM
TEL: +1-540-855-7041

ORDERS


SALES@GENUINBIOTECH.COM
FAX: +1-540-855-7041

WWW.GENUINBIOTECH.COM

4. Using a serological pipette, gently mix the lentiviral solution 3 times.
5. Carefully add 1 mL of the lentiviral solution to the well. Tip: To prevent splashing, add the solution to the dish along the wall.
6. Add a polybrene stock solution to the culture medium at a final concentration of 5 µg/mL. Gently swirl the dish to mix.
7. 48 h after cell release, without discarding the original medium, add another 1 mL of lentiviral medium directly into the dish.
8. Add an additional polybrene stock solution into the dish to obtain a final concentration of 5 µg/mL. Tip: Now, the medium in the dish should be a total of 3 mL.
9. 72 h after cell release, cells may reach confluence. Trypsinize the cells off the 35 mm dish and culture those cells in a 60 mm dish.
10. Add puromycin to the dish at a final concentration of 4 µg/mL. Tip: To assess the efficacy of puromycin selection, culture a dish of wild-type HeLa cells as a negative control.
11. Allow puromycin selection for 48 h. Almost all wild-type HeLa cells should die, while the dish infected with lentiviruses should have some remaining cells.
12. Replace the medium with regular growth medium without puromycin and allow the cells to grow to confluence before harvesting or staining.

Note: 1. This product is for research use only.
2. This product is only supplied to end users.
3. Do not use this product for commercial.

Validation Data



Genotype	Ct Value
Wild-Type	20.88
Knock-Down	22.03
$\Delta Ct (Ct_{KD} - Ct_{WT})$	1.15
% mRNA Reduction	↓ 55%

Copyright ©2024 Genuin Biotechnology LLC

RT-qPCR analysis. HeLa cells were infected with CDC16-specific shRNA lentiviral particles, total RNA was extracted from wild-type and knockdown cells, RT-qPCR was performed using gene-specific primers. $\Delta Ct (Ct_{KD} - Ct_{WT})$ was used to calculate mRNA reduction (%) between wild-type and knockdown cells using the following formula: $(1 - 1/2^{\Delta Ct}) \times 100\%$.

